

# Finance for the adaptation of ecosystems to climate change: a review of the Adaptation Fund portfolio

---

SEI working paper  
April 2019

Aaron Atteridge  
Sandra Tenggren





**Stockholm Environment Institute**

Linnégatan 87D 115 23 Stockholm, Sweden

Tel: +46 8 30 80 44 [www.sei.org](http://www.sei.org)

Author contact: Aaron Atteridge

[aaron.atteridge@sei.org](mailto:aaron.atteridge@sei.org)

Editing: Emily Yehle

Layout: Richard Clay

Cover photo: European bee-eater (*merops apiaster*), male with female, Italy

Photo credit: Tanja\_G / Getty

This publication may be reproduced in whole or in part and in any form for educational or non-profit purposes, without special permission from the copyright holder(s) provided acknowledgement of the source is made. No use of this publication may be made for resale or other commercial purpose, without the written permission of the copyright holder(s).

Second edition: Copyright © April 2019 by Stockholm Environment Institute

An earlier version of this working paper was published in December 2018. It was updated in April 2019 in response to feedback from the Adaptation Fund Board Secretariat. The Fund has limited scope to influence project priorities, and funded projects are defined by developing countries

The authors are grateful to the Adaptation Fund Board Secretariat for their helpful feedback, and to Nella Canales for her contribution to the analysis in this report

Stockholm Environment Institute is an international non-profit research and policy organisation that tackles environment and development challenges.

We connect science and decision-making to develop solutions for a sustainable future for all.

Our approach is highly collaborative: stakeholder involvement is at the heart of our efforts to build capacity, strengthen institutions, and equip partners for the long term.

Our work spans climate, water, air, and land-use issues, and integrates evidence and perspectives on governance, the economy, gender and human health.

Across our eight centres in Europe, Asia, Africa and the Americas, we engage with policy processes, development action and business practice throughout the world.

---

## Contents

<b>Abstract.....</b>	<b>4</b>
Keywords.....	4
<b>1. Introduction .....</b>	<b>5</b>
Protecting ecosystems and other species under climate change.....	5
The role of climate finance.....	6
Research outline.....	8
<b>2. Methodology .....</b>	<b>9</b>
<b>3. Findings: Is adaptation finance supporting ecosystems? .....</b>	<b>11</b>
How much finance is targeted at helping ecosystems deal with climate change? .....	11
Are ecosystem interventions emphasizing adaptation strategies or traditional conservation measures?.....	12
Is ecosystem resilience the main objective? .....	16
Exclusions .....	17
<b>4. Discussion .....</b>	<b>18</b>
More finance and more focus needed on the new challenges of managing ecosystems under climate change ..	18
Implications for climate funds and project developers.....	19
Concluding remarks .....	21
<b>References .....</b>	<b>22</b>

## Abstract

This paper is broadly interested in how much adaptation practitioners and funders are focusing on the problem of how species other than humans will adapt to climate change. Globally, human activities are decimating biodiversity, and climate change will magnify these threats and create new ones. At the same time, adaptation funds are limited; developing countries need far more funds than currently available, raising questions about how priorities are set and defined. There seems to be a high risk that, in a contest for limited resources, planners could ignore the plight of non-human species and instead prioritise actions that target specific human risks or vulnerabilities, thus leaving other species imperilled.

As an exploratory case study, we examine the project portfolio of one of the major international climate funds, the Adaptation Fund. The Fund has been given a mandate from the UNFCCC to respond to funding priorities identified by developing countries, which means countries identify the projects for funding. Our study provides insight into the kinds of projects being proposed, as well as the level of importance given — particularly by project developers — to the vulnerabilities and adaptation needs of non-human species.

Ecosystem outcomes are of limited importance to almost all projects, even those projects that include some ecosystem-related activities

First, we analyse the extent of funding that has been allocated for activities that address the needs of, and challenges faced by, other species. Second, we examine whether the activities funded were designed to reflect the priorities that conservation literature emphasises as necessary for building ecosystem resilience in the face of climate change, including the types of ecosystems prioritised and the scale of interventions. Third, we assess whether ecosystem resilience was the primary focus of the activities funded, or whether instead they were designed to use certain ecosystem elements to improve the resilience of people or reduce risks to people's physical and economic assets (sometimes referred to as "ecosystem-based adaptation"). Taken together, these sets of questions enable us to reflect on the level of urgency that countries — both donors and recipients — and adaptation planners are giving to the problem of how other species and ecosystems will cope with and adapt to climate change, and whether this problem needs more specific emphasis in the mandates and guidelines given to the climate funds.

We find that very little of the activities supported by the Adaptation Fund are ecosystem-related, totalling only about 15% of its allocated funding (as of late 2016). Of this, most funds go to activities that focus on delivering specific ecosystem "services" for people — such as protection against flooding or coastal erosion — rather than for other species. This likely diminishes any positive effects that projects are making towards ecological resilience. The most commonly targeted ecosystem types are forests, mainly through afforestation or reforestation activities; some of these are arguably of dubious benefit from an ecological perspective because they are re-planting exercises, rather than activities intended to re-establish complex ecological systems. None of the project proponents appear to have used regional or global ecological criteria as a basis for targeting particular ecosystems, nor do they explain in project documents the local ecological importance of the ecosystems. Indeed, most activities do not even address ecosystems as whole units but focus on individual components (e.g. planting of trees). On a geographic scale, most activities are miniscule. The conservation approaches employed are mainly traditional strategies rather than the kinds that conservation literature argues need greater emphasis in the face of climate change. Finally, the ecosystem-related outcome indicators included in projects' results frameworks are, with very few exceptions, only quantitative (e.g. number of hectares of land reforested), offering no evaluation of the quality of the outcomes from the perspective of particular species or overall ecosystem function. The latter reinforces our perception that ecosystem resilience is not a primary focus in most instances. Indeed, we estimate only around 4% of allocated funding has been directed to activities where ecosystem resilience is the primary objective.

We thus conclude that ecosystem outcomes are of limited importance to almost all projects, even those projects that include some ecosystem-related activities. Our findings raise serious questions about whether sufficient or meaningful attention and resourcing is given to how other species and natural ecosystems can adapt to climate change. We make recommendations on how climate funds, adaptation planners and project developers can address this.

## Keywords

Climate change, adaptation, ecosystems, finance

## 1. Introduction

### Protecting ecosystems and other species under climate change

Already under stress from myriad human activities, ecosystems globally are now increasingly threatened by climate change. This paper is broadly interested in how much climate funding is used to specifically address the plight of natural ecosystems (or individual species) in the face of climate change, and what kinds of interventions are being implemented.

Climate change will intensify the threats already facing non-human species, such as habitat degradation, landscape fragmentation, human-wildlife conflict, and the pollution of land, air and water. The pace of environmental change is now too fast for many species and ecosystems to adapt quickly enough (Jump and Penuelas 2005), and climate change could thus become the first or second largest driver of global biodiversity loss over the next decade (Sala 2000; Thomas et al. 2004). Poiani et al. (2011) anticipate the most common climate impacts will be changes to habitat quantity or quality and to hydrologic regimes. Other impacts include: sea-level rise; direct warming of habitats; increased fire frequency; pest outbreaks; increased spread of diseases, parasites and zoonoses; altered weather and precipitation patterns; glacial recession; increased populations of competitor species; and increased spread of invasive or non-native plants, animals and pathogens (Mawdsley et al. 2009). Further, as the environmental changes cascade, this is likely to create an unstable environment. For many ecosystems, this will increase the likelihood of damage and loss due to extreme events, which are expected to become more intense and more frequent in some regions (European Commission 2013; DEFRA 2008). A changing climate is shifting seasons and habitats, as well as changing life cycles and causing the emergence of new physical traits in species (Thomas et al. 2004; Millennium Ecosystem Assessment (Program) 2005); changing species distribution (Parmesan and Yohe 2003; Gimona et al. 2015; Pacifici et al. 2015; Mawdsley et al. 2009; Rüter et al. 2014; Burrows et al. 2014; Root et al. 2003; Gaston 2009); and changing migratory patterns (Cotton 2003; Otero et al. 2014; Singh et al. 2012). It is also leading to other effects, including the decoupling of co-evolved interactions like plant-pollinator relationships (Mawdsley et al. 2009). A swathe of species extinctions are predicted, in large part because ecosystem fragmentation and human land use means individual species will be unable to follow habitat range shifts (Urban 2015). Many vegetation types and individual species are expected to lose representation in protected areas (Araujo et al. 2004; Burns et al. 2003; Scott et al. 2002). Reserves at high latitudes and high elevations, on low-elevation islands and the coast, and with abrupt land-use boundaries are considered particularly vulnerable (Sala 2000; Shafer 1999). Current trends in biodiversity loss are doubly concerning in the face of climate change, because the capacity of ecosystems to cope with or adapt to environmental changes has been linked with having high levels of biodiversity and ecosystem heterogeneity (Vos et al. 2010; OECD 2003).

The rationales for addressing the plight of other species and natural ecosystems are rooted in morality (Brooke 2008; Thomas et al. 2004) but also in self-interest. Humans depend on other species and on different types of ecosystems for food, water, shelter, health, aesthetic and recreational value, cultural and spiritual value, and to buffer us against natural hazards including extreme storm events (Millennium Ecosystem Assessment (Program) 2005). Some estimate the economic losses associated with the loss of the world's natural capital to be between US\$ 2 million and US\$ 4.5 trillion (Feger and Pirard 2011). Studies on the accumulation of economic costs and benefits from the conversion of "wild nature" to human use landscapes have concluded that implementing an effective global program for conserving the remaining wild nature has a benefit-cost ratio of 100:1 (Balmford 2002).

There is a broad body of literature describing how, as a result of these new and intensifying threats, conservation practitioners may need to change the strategies they employ. The types of conservation strategies needed in the face of climate change have been described along a continuum, from promoting resistance, to enhancing resilience, and facilitating transitions (Millar et al. 2007; Glick, Stein and Edelson 2011; Glick, Chmura and Stein 2011). Resistance strategies attempt to maintain the status quo by minimising the exposure of species to climate impacts or by compensating for changes in the environment (such as by rebuilding habitat that might be degraded by climate change). *Resilience strategies* aim to enhance the ability of ecosystems or species to absorb or accommodate disturbances induced or exacerbated by climate change; in other words, the goal is to return ecosystems or species to a particular functional state, or to maintain some level of functionality despite being in an altered state (Holling 1973; Gunderson and Holling 2002; Heller and Zavaleta 2009). Transition strategies introduce the idea of

actively managing for a new functional state, for instance by preparing for changes in ecosystem types as coastal lands are inundated by rising sea levels, or by translocating species beyond current range limits in anticipation of future climatic conditions (Stein et al. 2013). There are expectations that, as climate change intensifies over time, resistance strategies may become more difficult to implement successfully (Millar et al. 2007), and conservation practices may need to shift from a focus on resilience towards managing for change (Chornesky et al. 2015; Lawler et al. 2015; Stein et al. 2013; West et al. 2009).

Various reviews of conservation literature have identified specific strategies that conservation might emphasise in light of climate change (Heller and Zavaleta 2009; Mawdsley et al. 2009; Hannah 2011). These strategies, as summarized in Table 1, include:

- protecting and managing large, intact landscapes and seascapes;
- protecting key geophysical features of the landscape;
- increasing the size and number of protected areas;
- maintaining and increasing landscape connectivity for species and ecological processes, by protecting movement corridors and managing the matrices between protected areas;
- identifying and protecting climate refugia (areas that are likely to become suitable habitats under future climatic conditions);
- ex-situ conservation, including establishing captive populations and gene banks;
- re-establishing species that are important ecosystem engineers; and
- reducing other non-climate related threats and stresses such as by tackling invasive species, pollution, and habitat loss due to human encroachment and land use change.

Adaptation policies and projects largely focus on how people's livelihoods and their physical and economic assets are in need of protection

More controversial strategies include translocating species, particularly those at risk of extinction, to more suitable habitats – as well as assisted evolution, as in the example of “super corals” (van Oppen et al. 2015).

In our analysis, we refer to these types of conservation strategies as “adaptation”, to denote their particular responsiveness to climate change (see “strategy type” in Table 1). These encompass the entire continuum mentioned above: promoting resistance, enhancing resilience, and facilitating transitions. They are distinguished in our descriptive analysis from what we refer to as “traditional conservation” approaches that focus either on restoration of degraded ecosystems or protection from current stresses such as pollution or human encroachment.

## The role of climate finance

At the international level, there is agreement in multiple forums on the need for urgent action to address these myriad threats to biodiversity, including in the Aichi Biodiversity Targets of the United Nations Framework Convention on Biological Diversity (CBD) (United Nations 1992a; Secretariat of the Convention on Biological Diversity 2012), and in various Sustainable Development Goals under the United Nations Agenda 2030 (such as goals 13 on climate change, 14 on conservation of oceans, and 15 on forests and biodiversity) (UN General Assembly 2015).

Addressing the needs of ecosystems and other species is also recognised as a key element of the climate adaptation agenda. The importance of accounting for climate change in the management of vulnerable ecosystems is explicitly mentioned in the strategic goals of the CBD (UNCBD 2016). Ecosystems are mentioned in Article 2 of the United Nations Framework Convention on Climate Change (UNFCCC), which explicitly states that countries need to reduce greenhouse gas emissions to the atmosphere “at a level that would prevent dangerous anthropogenic interference with the climate system... within a time frame sufficient to allow ecosystems to adapt naturally to climate change” (United Nations 1992b, p.4). This formulation seems increasingly outdated, however, as the conservation community has since been at pains to highlight that the rate of change is already so fast that many species are unlikely to be able to adapt “naturally”.

Yet in the evolving climate policy discourse, we observe that adaptation policies and projects largely focus on how people's livelihoods and their physical and economic assets are in need of protection. Far less emphasis is put on how to provide natural ecosystems and other species with targeted support to ensure their survival. This is evident even with increased attention to the concept of ecosystem-based adaptation

Table 1. Ecosystem strategies needing emphasis under climate change

Conservation parameter	Needs / approaches	References	Strategy type (as used in our analysis)
<b>Sites of conservation</b>	Focus not only on areas where target species occur today, but also areas with high potential for suitable habitats in the future (i.e. where species have high probabilities of persistence over the long term); identify important areas to protect in advance; create new protected areas.	(Hannah 2011; Donaldson et al. 2017; Oliver et al. 2012; Game, Lipsett-Moore, Saxon, et al. 2011; Ashcroft et al. 2009; Hodgson et al. 2009; Kareiva et al. 2008; Schmitz et al. 2015; Lawler et al. 2015; Nuñez et al. 2013; Hannah et al. 2007; Hannah 2015)	<b>ADAPTATION</b>
	Improve/maximise connectivity between habitat patches; facilitate movement of organisms, by building habitat networks, cross-environment connectivity, creating corridors across climatic gradients to enable range shifts, increasing landscape permeability to species movement, ecological networks; reduce fragmentation; conserve connectivity between climatically diverse areas.	(Heller and Zavaleta 2009; van Bodegom et al. 2013; Hannah 2011; Cross et al. 2012; Game, Lipsett-Moore, Saxon, et al. 2011; Hunter Jr. et al. 2010; Jones et al. 2016; Brooke 2008; Mawdsley et al. 2009; Poiani, Goldman, Hobson, et al. 2011; Ashcroft et al. 2009; Hodgson et al. 2009; Vos et al. 2008; Environmental Defenders Office 2009; Heller et al. 2015; Hannah et al. 2008; Wilson and Piper 2008; Theobald et al. 2015; Groves et al. 2012; Morecroft et al. 2012; Krosby et al. 2010; Hulme 2005; Kareiva et al. 2008; Bonn et al. 2014; Tischendorf and Fahrig 2000; Schmitz et al. 2015; Stein et al. 2013; Lawler et al. 2015; Hannah et al. 2002; Vos et al. 2010; Nuñez et al. 2013; Halpin 1997; Noss 2001; Lawton et al. 2010; Alagador et al. 2016; Johnson and Welch 2009; Settele et al. 2014; Townsend and Masters 2015)	
<b>Scale</b>	Shift from site-scale to landscape-scale conservation, protect and manage large landscapes; expand remit from nature reserves and patch-based management to the wider landscape.	(Hannah et al. 2008; Heller and Zavaleta 2009; Clews 2012; van Bodegom et al. 2013; Vos et al. 2010; Stein et al. 2013; Theobald et al. 2015; Ackerly et al. 2010; Mawdsley et al. 2009; 2017; Lindenmayer et al. 2007; Lawton et al. 2010)	
<b>Choice of species and managing diversity</b>	Representation: protecting a portfolio of variant forms of a species or ecosystem so that, regardless of the climatic changes that occur, there will be areas that survive and provide a source for recovery; focus on maintaining diversity, structure and function, rather than attempt to preserve current species composition; rethink the mix of species to be planted and potentially focus on ecosystem function rather than particular assemblages of species; manage keystone species or species at risk of extinction; increase heterogeneity of landscapes.	(Hannah et al. 2008; Harris et al. 2015; Starzomski 2013; Mawdsley et al. 2009; Shoo et al. 2013)	
	Emphasise high level of biodiversity, as an important prerequisite for the adaptive capacity of ecosystems	(Vos et al. 2010; 1996; Hopper 2007; Brooke 2008)	
<b>Translocation</b>	Assisted migration, assisted colonisation, translocation of species that will not survive in situ	(Harris et al. 2006; Hoegh-Guldberg et al. 2008; Thomas 2011; Hole, Huntley, Arinaitwe, et al. 2011; Hannah 2011; Environmental Defenders Office 2009; Clews 2012; Gallagher et al. 2015; Lunt et al. 2013; Poiani, Goldman, Hobson, et al. 2011; Mawdsley et al. 2009; Hulme 2005; Kareiva et al. 2008; Krosby et al. 2010; Stein et al. 2013; Lawler et al. 2015; Hannah et al. 2007; McClanahan et al. 2008; Pearson and Dawson 2005)	
	"Ex situ" conservation: gene and seed banks, establishment of captive populations of species at risk of extinction, captive breeding	(Heller and Zavaleta 2009; Mawdsley et al. 2009; Hannah 2011; Tingley et al. 2014; Hunter Jr. et al. 2010; Settele et al. 2014; Department of Environment, Climate Change and Water 2010; Stein et al. 2013)	
<b>Fostering or working with ecosystem change</b>	Convert one type of ecosystem to another; prepare an ecosystem to function differently in a future climate, rather than restoring a historical condition	(Wilson and Piper 2008; Keppel et al. 2012; Game, Lipsett-Moore, Saxon, et al. 2011)	
	Establishment of climate refugia: conserving representative examples of geophysical settings will protect representative ecological communities under both current and future climates	(Beier and Brost 2010; Stein et al. 2013; Mawdsley et al. 2009; Theobald et al. 2015; Jones et al. 2016; Game, Lipsett-Moore, Saxon, et al. 2011; Morecroft et al. 2012; Groves et al. 2012; Schmitz et al. 2015)	
<b>Managing for extreme events</b>	Measures to diminish the likely impacts of extreme storm events on ecosystems	(Bonn et al. 2014; European Commission 2013)	
<b>"Traditional" conservation</b>	Ecosystem protection: Reducing existing threats not related to climate change (e.g. land use change, fragmentation of habitat, pollution, human conflict, invasive species)	(Poiani, Goldman, Hobson, et al. 2011; Mawdsley et al. 2009; Hulme 2005; Kareiva et al. 2008; Krosby et al. 2010; European Commission 2013; Convention on Biological Diversity 2016; Bonn et al. 2014; Morecroft et al. 2012; Stein et al. 2013)	<b>PROTECTION</b>
	Habitat restoration or rehabilitation	(Maciver and Wheaton 2005; Harris et al. 2006; Millar et al. 2007; Environmental Defenders Office 2009; Clews 2012; Poiani, Goldman, Hobson, et al. 2011; Mawdsley et al. 2009; Hulme 2005; Kareiva et al. 2008; Krosby et al. 2010)	<b>RESTORATION</b>

(EBA), where the main focus is ensuring the resilience of people rather than other species (Martin 2011). While EBA projects can materially improve the prospects for some other species, EBA proponents usually pick and choose which ecosystem “services” to strengthen on the basis of targeted outcomes for *people*, while ignoring other elements that may be critical to broader ecological health.

As part of the global response to climate change, it is critical that as project developers identify and design adaptation projects – and conduct dialogue with funders – they give explicit attention to the needs and vulnerabilities of other species, and of whole ecosystems. This does not mean every project needs to focus on ecosystems. However, we argue that it does mean that adaptation portfolios need to ensure sufficient support is given specifically to addressing the vulnerabilities and adaptation needs of other species.

Large volumes of finance are needed to implement national biodiversity strategies and the objectives of the CBD, as well as to foster the mainstreaming of biodiversity considerations into other sectors like agriculture and trade (Richerzhagen et al. 2016) and to support adaptation in the face of new threats arising from climate change. There are significant methodological challenges in trying to assess how much funding is needed to achieve international goals (Feger and Pirard 2011). In one estimate, McCarthy et al. (2012) suggest that conservation funding needs to increase by an order of magnitude over present levels. In 2014, Parties to the CBD agreed to double financial resources to biodiversity protection, to flow particularly to the least developed countries, small island developing states, and economies in transition (UNCBD 2014). There have also been calls to increase finance for conservation and biodiversity in the Addis Ababa Action Agenda, in the 2015 Finance for Development Conference, and in Agenda 2030 to promote the Sustainable Development Goals.

The burden for financing ecosystem protection and adaptation does not fall solely on the actors involved in programming “climate finance”.<sup>1</sup> However, supporting ecosystems is a critical part of the global response to climate change. The UNFCCC and other institutions that support developing countries in addressing climate change — such as the World Bank, which governs several of its own Climate Investment Funds — need to ensure sufficient support for improving the resilience and adaptive capacity of diverse ecosystems and other species. There are few estimates of the likely costs of achieving biodiversity protection specifically in the context of climate change, including adaptation needs. Those that exist are mostly at the level of individual species in particular countries. A 2007 report to the UNFCCC (Berry 2007) estimates that the adaptation of natural ecosystems to climate change could cost US\$ 385 billion per year globally, under a business-as-usual emissions scenario and including further investments to expand terrestrial and marine protected areas by 10%. However, such figures are highly uncertain. The UNFCCC report also is now more than a decade old; actual financial needs may have since risen.

Under the UNFCCC, Parties have committed to mobilize financial support specifically to help developing countries tackle the impacts of climate change. Many of the world’s most biodiverse and ecologically threatened regions are in developing countries, and many developing countries are highly dependent on international financial support, so how these flows of climate finance are used is likely to have a crucial bearing on the resilience of ecosystems and other species to the impacts of climate change.

## Research outline

To date, there has been little analysis of whether adequate funding is being mobilized for global conservation efforts broadly (Hannah 2011), or, more specifically, how extensively the funding being mobilized for climate change is supporting the adaptation needs of non-human species.

In this paper, therefore, we examine the scale and character of financial support given to ecosystems through the Adaptation Fund. The Fund, set up under the UNFCCC, began financing projects in 2010. It is not the largest climate fund in operation, but was selected for this exploratory case study because it is the first major fund set up specifically to support adaptation to the impacts of climate change. It also has a sufficient and manageable project portfolio for detailed examination. The Adaptation Fund receives its

<sup>1</sup> Climate finance is not a clearly defined concept. Usually it refers to the financial commitments that countries under the UNFCCC have committed to provide to developing countries for the purposes of tackling both mitigation of greenhouse gas emissions and adapting to the impacts of climate change. It consists of funds mobilized by “donor” countries and channeled either bilaterally to developing countries or through multilateral development finance institutions and funds.



guidance from the UNFCCC process, and the core of its mandate is to support adaptation programmes and projects that are country-driven. This means projects funded reflect prioritisation decisions by developing countries themselves, as opposed to being set within the Fund. Although its Results Framework defines one of its seven strategic outcomes as “increased ecosystem resilience in response to climate change and variability induced stress” (Adaptation Fund 2011), the Framework is not used to dictate funding priorities for developing countries. Instead, it is used to report *ex post* on how projects approved by the Fund are contributing to the achievement of the Fund’s overall goal and outcomes. In other words, the projects funded by the Adaptation Fund to date are defined by developing countries, and the Fund itself has limited scope to influence the prioritisation process. However, the Fund’s Secretariat conducts technical reviews of proposals and has scope to provide feedback to project developers, to ensure the best possible outcomes are achieved with its limited funding. Hence, through dialogue, the Fund can help shape project activities.

In our analysis, we first calculate how much of the support provided by the Adaptation Fund has been used for ecosystem-related activities. Second, we examine whether the funded ecosystem activities reflect, in design, the strategies prioritised by conservation literature for building ecosystem resilience to climate change (as opposed to applying traditional conservation measures). Specifically, we examine: the type of ecosystems targeted, the physical scale of the interventions, and the type of conservation strategy. Third, we look at whether ecosystem outcomes are in fact important to the primary logic or objective of the projects. In other words, we ask: Is ecosystem resilience the main objective of the activities?

By answering these questions, we are able to reflect more deeply about whether adaptation planners, funders and the UNFCCC itself are giving sufficient attention to the plight of ecosystems in the face of climate change.

## 2. Methodology

As of September 2016, the Adaptation Fund had approved support for 53 projects. The first step in our analysis is identifying ecosystem-related activities within these 53 projects and delineating their budgets.

First, we review all of the relevant project approval documents available on the Adaptation Fund’s website.<sup>2</sup> For each project, we classify as “ecosystem-related” any activities that (i) directly intervene with one or more ecosystem elements (or focus on a particular species) and (ii) suggest an ambition to materially improve the health or functioning of these ecosystems (or species). Our focus is on *natural ecosystems*, which we acknowledge is a problematic concept to clearly define. The main distinction we make is between landscapes that are not under intensive human use (“natural ecosystems”) and those that are (e.g. agricultural or urban biomes). For highly disturbed landscapes, such as some grasslands included in several projects, we include it as a “natural ecosystem” if the intention of project activities is to recover the natural grasslands ecosystem (in other words, if it is designed from the perspective of non-livestock species). However, the project is excluded if it rehabilitates grasslands to improve livestock grazing prospects.

We then extract budgets for these activities, to the extent these are delineated in overall project budgets. We exclude general project management fees that are often included in project budgets. Although there are costs to be borne when implementing projects, these are only reported in project documents on a lump-sum basis, rather than broken down into individual activities; they also are related to the transaction costs of project-based funding modalities, rather than the costs of undertaking the ecosystem-related activities.

Our next step is to qualitatively examine these activities by reviewing the project-, component- and activity-level descriptions, in order to answer the research questions related to the design of ecosystem-related activities. Again using the project documents, we manually code the following information about each of the ecosystem-related activities:

By answering these questions, we are able to reflect more deeply about whether adaptation planners, funders and the UNFCCC itself are giving sufficient attention to the plight of ecosystems in the face of climate change.

<sup>2</sup> <https://www.adaptation-fund.org/>

- i. what kinds of ecosystems are targeted, noting not only physical type but also any indications of whether the ecosystem was selected on the basis of any ecological criteria;
- ii. what physical scale the intervention has; and
- iii. what specific conservation strategies are proposed.


We classify the latter strategies (iii) against the spectrum of conservation strategies described in Section 1, namely whether they explicitly include any measures that match those given emphasis by conservation literature as important in the face of climate change (what we refer to as “adaptation” measures), or whether they adopt more traditional restoration or protection measures.

We classify as “adaptation” strategies those listed in the literature summary in Section 1. These include, for example, improving connectivity between ecosystem fragments and enabling migration, protecting climate refugia, and protecting key geophysical landscape features. “Restoration” strategies are activities designed primarily to rehabilitate or restore ecosystems that have been degraded or destroyed, such as through reforestation or tackling pollution sources. “Protection” strategies are those that aim to reduce current stresses on ecosystems, for instance by establishing conservation areas.

Finally, to address our third research question we manually code from project descriptions how the narrative relates to ecosystems versus other potential project beneficiaries. We also code from project monitoring and evaluation (M&E) frameworks what specific indicators of ecosystem *outcomes* have been included.

Figure 1 summarises our methodology and explains the link with each of the research questions.

Figure 1. Methodology for analysing the Adaptation Fund portfolio



Overall, our findings probably overestimate the number of ecosystem-related activities that have been supported by the Adaptation Fund, and the budgets for these. We give the benefit of the doubt and include some projects where the relevance to ecosystems is inferred but not entirely clear. Project budgets are not always sufficiently disaggregated to allow us to isolate ecosystem-related activities from other project activities, and sometimes budgets combine several different activities together. Where we are unable to delineate budgets for ecosystem-related activities specifically, we include the larger component budgets in our finance estimates and thus overestimate the level of financial support.

### 3. Findings: Is adaptation finance supporting ecosystems?

#### How much finance is targeted at helping ecosystems deal with climate change?

The total volume of finance committed by the Adaptation Fund to its first 53 projects in developing countries is US\$ 346.2 million. Commitments for individual projects range from US\$ 690,000 up to US\$ 10 million, with an average of around US\$ 6.5 million per project.

Across this portfolio, we identify ecosystem-related activities in 23 projects, or 43% of the total projects funded. These are summarized in Table 2. In most cases, these activities are not the primary focus but are a relatively small part of a larger project. Their aggregated budget (US\$ 52.3 million) accounts for approximately 15% of total Adaptation Fund commitments. Note that the finance figures represent approved amounts rather than actual disbursed expenditure.

**Table 2. Summary of ecosystem-related activities within Adaptation Fund projects**

Project country	Ecosystem-related activities	Budget of ecosystem related activities (US\$)	Total project budget (US\$)	Ecosystem type(s)
Belize	Component 1: Improving protection regime of marine and coastal ecosystems; Component 2: Alternative livelihoods to reduce vulnerability to reef decay and/or protect the reef from further degradation; Component 3: Awareness raising and building local capacity.	4 600 000	6 000 000	Coral reef, mangroves
Cambodia	Output 1.3: Forest restoration protocols developed for CPA intervention sites; Output 2.2: Restoration of degraded forest to multi-use forest.	1 648 125	4 954 273	Forest
Colombia	Component 2: Rehabilitation of wetlands and their hydrology in the target area as a means to reduce risk to flooding and drought associated with climate change and variability; Specifically, Output 2.2: rehabilitation of ecosystems associated with hydrodynamics of the target areas.	1 459 530	8 518 307	Wetlands
Costa Rica	Components 2.1 and 2.3 include relevant activities. Activity 2.2.1 indicates (for example) "reforestation at aquifer recharge areas". Output 2.3, Activities 2.3.1, 2.3.2 and 2.3.3 involve coastal environment management plans and mangrove replanting and conservation.	1 114 724	9 970 000	Mangroves
Cuba	Component 1: Recovery of coastal ecosystems; Components 2 and 3: Participatory management and enabling environments relating to ecosystem-based adaptation (EBA).	5 220 000	6 067 320	Mangroves, coastal wetlands
Ecuador	Output 2.1.3: Natural resource assets created, improved or maintained	1 525 000	7 449 468	Forest
Honduras	Output 2.1: Enhance connectivity in protected forests	155 000	5 620 300	Forest
India (Andhra Pradesh)	Outcome 3: Restore mangroves to overcome salinisation and other impacts due to sea level rise, including nursery establishment.	106 950	689 264	Mangroves
Indonesia	Outcome 1.2: Village conservation agreements; Outcome 2.3 reforestation and afforestation.	458 600	5 995 666	Forest
Jamaica	Component 1: Install hard engineering breakwaters, allowing regeneration of seagrass beds.	5 480 780	9 995 000	Seagrass beds
Kenya	Component 3: Integrated shoreline and mangrove ecosystem management; Outputs 3.2 (mangroves) and 3.3 (coral reefs)	703 916	9 998 302	Coral reef, mangroves
Lebanon	Output 3.1: Community-based sustainable rangeland management plan; Output 3.2: Restore degraded rangeland areas and reduce flood risks.	2 550 000	7 860 825	Grasslands
Madagascar	Output 2.2: Watershed rehabilitation including reforestation (also includes other activities not related to ecosystems);	460 000	5 104 925	Forest
Mauritius	Output 1: Includes hard engineering structures that will improve ecosystem outcomes (new habitat, improved biodiversity compared with baseline); 1.3 includes mangrove plantation.	4 870 000	9 119 240	Mangroves, coastal ecosystems
Mongolia	Component 1: Landscape-level integrated land use and water resources monitoring and planning system; Output 2.1: Local level climate change adaptation assessment and monitoring; Output 2.2: Integrated landscape level, ecosystem-based adaptation management action plans; Output 2.3: Physical techniques to improve ecosystem resilience; Output 2.4; Component 3: Institutional and policy capacity strengthened to support ecosystem-based adaptation replication, monitoring, and enforcement for critical watersheds.	4 589 124	5 500 000	Forest

Project country	Ecosystem-related activities	Budget of ecosystem related activities (US\$)	Total project budget (US\$)	Ecosystem type(s)
Myanmar	Output 1.2: Micro-watersheds protected and rehabilitated through Farmer-Managed Natural Regeneration (FMNR) to increase natural water retention and reduce erosion. Relevant activities include 1.2.5, 1.2.6, and 1.2.7, as well as capacity building and preparatory work.	1 374 000	7 909 026	Forest
Nicaragua	Component 2: Sub-activity on protecting small area of forests around recharge areas and riparian zones.	100 000	5 500 950	Forest
Papua New Guinea	Output 1.3: Support system for community-led mangrove reforestation and conservation projects.	455 000	6 530 373	Mangroves
Peru	Component 1: Introduction of sustainable fishing methods; Restoration and co-management of natural banks; Introduction of sustainable aquaculture; production of bio-fertilisers from fishery and aquaculture residue (reducing marine pollution); Component 2: Coastal marine environmental surveillance; Component 3: Capacity building and knowledge management for EBA; Component 4: Management and regulation for coastal ecosystem resilience.	5 850 000	6 950 239	Marine ecosystems (fisheries)
Seychelles	Component 1 Reforestation and removal of invasive species; Component 2 Rehabilitation of tidal wetlands and coral reefs; Component 3 Policy framework and training in EBA.	5 495 000	6 455 750	Forest, wetlands, coral reefs
South Africa (KwaZulu-Natal)	Component 2: Climate-proof settlements through "ecological and engineering solutions"; Output 2.2: Restored and protected critical ecosystems that maintain ecosystem resilience, provide buffering from climate change impacts and provide freshwater to local communities downstream. Activity 2.2.1: Restore and rehabilitate critical ecological infrastructure to improve its capacity to mitigate effects of climate induced disasters.	823 810	7 495 055	Grasslands, wetlands
Tanzania	Component 2: Coastal ecosystems rehabilitated and Integrated Coastal Area Management implemented; Introduces efficient cookstoves to reduce fuelwood demand and effects on mangrove forests; Component 3 includes as one output: "One Ecosystem Based Integrated Area Management (EBICAM) plan for the coastal region approved".	479 000	5 008 564	Coral reef, mangroves
Uganda	Activity 2.1.1: Identifying most degraded forest areas vulnerable to rainfall; Activity 2.1.2: Afforestation; Activity 2.1.4: Rehabilitation of degraded wetlands; Activity 2.1.5: Restoration of degraded riverbanks.	2 828 500	7 751 000	Forest, wetlands
<b>Total</b>		<b>52 347 059</b>	<b>156 443 847</b>	

As illustrated in Figure 2, projects with ecosystem-related activities allocate a relatively small portion of the project's total financial commitment to such activities; this allocation averages 33% for the 23 projects. In seven projects, the relevant budget is more than 50% of the total project budget, while in another seven projects the allocation for ecosystems makes up less than 10% of the total budget. The average amount of funding for ecosystem-related activities was US\$ 2.3 million. Four projects include a budget of more than US\$ 5 million, the largest amount being in Peru (US\$ 5.58 million). At the other end of the scale, nine projects budgeted less than US\$ 1 million for ecosystem activities.

In at least nine projects, some of the activities we categorize as ecosystem-related focus on capacity building or the strengthening of legal or policy frameworks, rather than on implementing conservation activities on the ground. In several cases, such as Mongolia, these are the main relevant actions of the project.


## Are ecosystem interventions emphasizing adaptation strategies or traditional conservation measures?

### Types of ecosystems or species that projects are targeting

The ecosystem-related activities identified in Adaptation Fund projects focus on various ecosystems. As shown in Figure 3, the most common are forests, which are included in 10 different projects. These are mostly reforestation or afforestation activities. However, the ecosystems are almost always described in vague terms and usually without specifying forest type or quality, the mix of trees to be planted, the ecological outcomes for different species, how the land will be used or managed after being reforested, or any clear outcomes for biodiversity or the forest ecosystems themselves. Based on the language of project documents, many of these activities are probably actually tree plantations – often “multi-use” trees



Figure 3. Types of ecosystems funded in projects.



Note: Several projects include activities that target more than one ecosystem type, in which case both ecosystems are included here. This explains why the total number of ecosystems categorised (32) exceeds the total number of projects (23). It is not possible to separate total funding by ecosystem, since several projects intervene in more than one ecosystem type but usually do not break down the project budgets for different ecosystems separately.

Based on project documents, we see no evidence that the selection of ecosystems was based on any ecological criteria, such as the presence of threatened or iconic species, the area’s vulnerability to climate change, or the locations of biodiversity hotspots. None of the projects propose activities in sites of declared ecological importance (e.g. home to endangered species) or under current protection status. Almost all the environments targeted by these projects have previously been significantly degraded.

**What is the physical scale of ecosystem activities?**

Also noteworthy is that the scale of ecosystem interventions is sometimes very small in area. For example, in Nicaragua, forest restoration efforts target 50 hectares around water recharge areas. In Jamaica, major engineering works aim to rehabilitate nearshore seagrass beds along a one-kilometre stretch. Such small interventions have a correspondingly small benefit for biodiversity, and in some cases it is questionable whether there is any significant benefit at all. Some activities do cover larger areas – such as those in Cuba (about 7,300 hectares of mangroves), Colombia (700 hectares of wetlands) and Myanmar (4,200 hectares of micro-watersheds) – but the relevant activities tend to cover only part of the specified areas, and it is not always clear how much will actually be restored.

Almost none of the projects describe how the particular activities will fit into the wider landscape, such as whether they link up separate fragments of intact natural habitat (Honduras is an exception here).


### What kind of conservation approaches are being used?

As mentioned in Section 1, a swathe of conservation literature has highlighted the kinds of conservation strategies that will likely need more emphasis in the context of climate change. Here, we examine whether the ecosystem-related activities supported by the Adaptation Fund reflect these recommendations in their design (what we refer to in shorthand as “adaptation”), or whether they adopt a more traditional mix of conservation strategies (what we refer to as “restoration” and “protection”).

As shown in Figure 4, we identify only one project, in Honduras, that includes activities in our “adaptation” category. This project document describes a focus on establishing connectivity in forest ecosystems. It is, however, a very small component (US\$ 155,000) of a much larger project, and the larger project includes no other ecosystem-related activities, so the likely impacts are small. Two other projects reference “landscape” approaches that align with our adaptation category. The Seychelles project document specifically mentions that the project aims to improve landscape connectivity, between different ecosystem types and within the same ecosystem type. However, since we are not able to identify which activities achieve this in the project description or budget, the relevant project activities are categorised predominantly in the “restoration” category. The Uzbekistan project document mentions the adoption of a landscape-wide approach and discusses the interconnectedness of different parts of the patchwork landscape. However, the target landscapes are and will continue to be under intensive human use, meaning we identified no ecosystem-related activities within the project.

Most activities (19) focus on restoration to improve the condition of degraded ecosystems. Relatively few (6) are designed primarily to protect ecosystems from current stresses like pollution. Five projects – in Belize, Indonesia, Seychelles, Tanzania, and Uganda – contain elements of both restoration and protection, and in Figure 4 we have delineated budgets to separately categorise each strategy (which explains why there are 28 coded results for 23 projects). In some other cases, there are suggestions of both types of

Figure 4. Conservation approaches in projects with ecosystem-related activities.



activities within an overall project. However, the delineation between these is not clear and/or separate budgets cannot be extracted from project documents; in such cases, the entire project was categorised in Figure 4 according to the larger strategy.

Overall, we find approximately US\$ 41 million in funding is committed to restoration activities (equivalent to 12% of the Adaptation Fund's total portfolio), and US\$ 11.2 million is committed to protection activities (3% of the total portfolio). Only a very minor amount of US\$ 155,000 supports the "adaptation" category.

### Is ecosystem resilience the main objective?

The inclusion of ecosystem-related activities does not necessarily mean the activities were designed specifically to improve the resilience of ecosystems or other species. Figure 5 shows the breakdown of funding across the whole Adaptation Fund portfolio, between (i) activities that are designed to address ecosystems specifically, (ii) activities that intervene in ecosystems but are designed mainly to deliver resilience outcomes for people or infrastructure, and (iii) activities that are not related to ecosystems. Just US\$ 15 million, or 4% of the total Adaptation Fund portfolio, is designed primarily for ecosystem resilience. This amounts to less than one third of the ecosystem-related budgets.


The budget targeting ecosystem outcomes consists of the funding for projects in Belize, Peru and Mongolia. The Belize project aims to strengthen the climate resilience of coral reefs; improving ecological health is stressed as a key purpose of the overall project. The Peru project targets two marine ecosystems, and highlights climate (and other) risks to these, and their importance as fisheries. The activities in Peru also clearly aim to provide benefits for people, in terms of improved fisheries resources; however, since the project document identifies specific target species, and biodiversity, as key project outcomes, we determine ecosystem outcomes to be a clear purpose and focus. The Mongolia case is less clear but does specifically describe habitat restoration in its results framework, and we give it the benefit of the doubt.

Across the remaining projects with ecosystem-related activities, the focus – and hence choice of ecosystem and design of intervention – is on increasing resilience of vulnerable people rather than on specific ecological outcomes. This includes projects that use the term "ecosystem-based adaptation" but typically focus on the provision of benefits for human well-being while providing little or no explanation of how other species will benefit from the intervention. In such cases, it is not clear whether other species do benefit – or might even be made worse off – because the description of the local ecosystem and how it functions is superficial in project documents.

A further means of examining whether ecosystems are central to a project's logic is to look at how the intended ecosystem *outcomes* have been defined by proponents in project monitoring and evaluation frameworks. Only two of the 23 projects we classified as having relevant activities include any indicators that are related to the quality of the ecosystem, such as to the resilience or survivability of species. The Cuba project proposes to develop a range of indices, including some that may help to gauge the state of the ecosystem (e.g. canopy density). The Peru project includes the design of an ecosystem resilience monitoring program, which is not in itself an indicator of ecosystem quality but at least suggests ecosystem health might be monitored. All other projects include indicators based only on numbers of activities or on physical areas (e.g. the size of the area of mangroves planted or of coral reef rehabilitated or of forests placed under conservation agreement). The India project includes an indicator related to growth and survival rate of trees, which is not a measure of the ecosystem per se but at least is one step advanced from simply measuring the number of hectares planted. The Jamaica project, which we classified as designating a large budget towards ecosystem-related activities, in fact had no indicators at all relating to ecosystem outcomes. Overall, based on the projects' results frameworks, we conclude that ecosystem outcomes are actually of limited importance to almost all projects.



**Figure 5. Value of Adaptation Fund commitments according to intended main beneficiaries (i.e. ecosystems or people)**



## Exclusions

Many project documents make at least passing references to benefits for ecosystems, but rather fewer actually describe and allocate funding for activities that might materially improve the health and resilience of ecosystems. Overall, while we are generous in our decision of which activities to include or exclude as ecosystem-related (meaning we tended to include rather than exclude), we exclude some projects that refer to ecosystems in their project documents but did not seem to include tangible activities. An example is the Cook Islands; the project document describes conservation of coastal, inland and reef ecosystems under the heading “environmental benefits” but does not describe any ecosystem-related activities in its work plan nor contain any monitoring and evaluation indicators that relate to these outcomes.

Several projects are particularly difficult to classify. The Guatemala project document clearly describes the impacts climate change will have on natural ecosystems in the project areas, and the specified objective of one of its four components is “development and implementation of climate change resilient ecosystem management”. However, closer review of the activities show that they focus on introducing agro-silvo-pastoral practices, by planting native trees and protecting soil degradation in cropping areas. A small subset of activities is listed as “climate resilient forest restoration activities that are more suited to high risk areas – berms, bunds, terraces, gully plugs, etc. – in order to increase landscape resilience as a whole under emerging long-term climate conditions. This includes lands designated for reforestation, conservation of existing ecosystems (and their ecosystem services), and promotion of small forest enterprises.” However, from the detailed description of project activities, we cannot ascertain any clear investments in ecosystems. The Uruguay project document describes the expected results in terms of protection and restoration of natural grassland biodiversity and talks about enhancing ecosystem services – mainly water provision for farmers and livestock – but then does not specify how this will be

achieved in practice, other than through “adaptation interventions”. Moreover, the interventions target the maximisation of livestock productivity on the grasslands, while no clear biodiversity outcomes are defined and there seem to be no measures that are likely to improve the grasslands ecosystem from an ecological perspective.

Many projects have budgets for vegetation planting, referred to under various names including reforestation or afforestation. Vegetation planting by itself is not automatically classified as ecosystem-related; we include it only when the project document describes some semblance of a natural ecosystem or refers to addressing the needs of specific species. As mentioned, virtually none of the project documents define how reforestation activities will strengthen the resilience of particular species or ecosystems, how ecosystem functions and structures are being restored, or even whether the tree species being planted are native. Among those activities that we do not consider as ecosystem-related are those focusing on the planting of multi-purpose tree species for human use and harvesting (Ghana, India, Mali) and the planting of “vegetative buffers” along coasts or floodways that are essentially a thin strip of trees (Georgia). Several projects – in Uzbekistan, Mauritania, Pakistan, Rwanda and Nepal – include the planting of trees and shrubs to reduce soil erosion but do not mention any natural ecosystem; these activities do not describe whether this stabilisation work improves or disturbs the natural ecosystem, nor how it changes conditions for other species. Consequently, we also do not consider these activities as ecosystem-related.

A number of projects are structured around community-based activities and/or the use of small grants, where part of the project itself is to identify and plan activities with local communities. For example, the Cook Islands project may contain small grants for community programs, and similarly, the Small Grants Facility of the South African National Biodiversity Institute (SANBI) mentions “ecosystem resilience.” But both are at the project approval stage and have not yet defined specific activities. In these cases, it is not possible for us to identify specific activities that are targeting ecosystems, even if the proposals mentioned that activities might eventually include reforestation or other relevant activities.

## 4. Discussion

### More finance and more focus needed on the new challenges of managing ecosystems under climate change

Our analysis reveals that natural ecosystems do not appear to be receiving much attention in the projects developing countries bring forward to the Adaptation Fund. Having reviewed only one of the international funds, it is not possible to say whether this is part of a broader pattern. However, we are concerned that it might be, given that other climate funds, like the Green Climate Fund, were set up with a similar country-driven mandate.

Overall, the *scale* and *quality* of interventions that we classified as ecosystem-related appear highly questionable from the perspective of addressing the needs and vulnerabilities of other species. Even with our generous approach to classification, only 15% of total finance provided through the Adaptation Fund has been for ecosystem-related activities and only 4% has specifically targeted ecosystem needs. These amounts are spread into mostly small activities across different projects, with very few large, dedicated interventions – either in physical size or scope of financial support. The physical area of ecosystems included is sometimes miniscule in size, such that it is difficult to imagine any meaningful impact on overall ecosystem resilience or the survivability of other species. This contrasts with the recommendations of conservation literature to move beyond small conservation efforts and focus instead on landscape-scale interventions (see Table 1). The results also highlight that high conservation value ecosystems are not included in projects, as is recommended in conservation literature. Instead, most of the activities propose to work within highly degraded environments. This perhaps explains why we classified almost all of the activities as “restoration”, and only one very small activity as “adaptation” targeting natural ecosystems.

Even though project documents often allude to ecosystem benefits, for the most part these are not reflected in monitoring and evaluation frameworks, leading us to conclude they are not important to the project’s purpose or logic. As a result, it is possible that even among the project activities we classified as ecosystem-related, in reality many may deliver little or no tangible benefit to other species.

Even among those projects that include ecosystem-related activities, most have not been designed with the resilience of ecosystems in mind

Indeed, even among those projects that included ecosystem-related activities, most have not been designed with the resilience of ecosystems in mind; we estimate only 4% of total funding commitments from the Adaptation Fund were directed to activities specifically targeting ecosystem resilience. Instead, projects include specific ecosystem components in order to improve the resilience of vulnerable people or reduce risks to people's physical and economic assets. This does not mean the activities generate no benefits for ecosystems, but the selection of what to do and how to approach the conservation effort was likely not made based on the needs of other species. An example here are some of the "reforestation" activities, which appear to be designed as tree planting exercises rather than ecological restoration.

Going further, projects usually define ecosystems in vague and simplistic terms. Project documents tend to describe only generic features of an ecosystem, if any at all (many forest-related projects do not), and leave ecological characteristics poorly defined. There is rarely discussion of specific species, or the inter-relationship between species, or of important ecosystem properties, as opposed to ecosystem "services". Most of the ecosystems included in projects are significantly degraded, and none were selected by proponents on the basis of high ecological merit – though we recognize that some, such as coral reefs or wetlands, may in fact be ecologically important in a local sense. Project documents often make no explicit statements about what kinds of ecosystem outcomes might be expected, in terms of biodiversity, population size, habitat heterogeneity, or changes relative to the current condition of the landscape. This in itself suggests that ecosystem outcomes are not being treated with great importance by project developers.

Most of the project documents provide no clear description of the suitability of the chosen interventions, their likely effectiveness (*vis-à-vis* alternative strategies, for example), what changes they will actually produce for biodiversity, or what contribution they might make to conservation goals or ecosystem adaptation.

With one very small exception, our results also suggest that ecosystem interventions are not applying strategies that reflect those emphasised in conservation literature as critical for protecting other species in the face of climate change. Most activities we identified involve traditional restoration strategies at a very local scale and are not contextualised in the wider landscape; they have not been designed to maximise conservation or biodiversity outcomes or to address the future vulnerabilities and needs of other species as the impacts of climate change intensify. This may be partly related to the fact, highlighted above, that ecosystems are rarely the main intended beneficiaries of the interventions. Most instead seek to generate discrete ecosystem "services", like soil stability, water provision, or protection against coastal erosion and storm surges.

In practice, conservation of ecosystems and individual species will need to work with a mix of these "restoration", "protection" and "adaptation" strategies. What our findings suggest is that at present, adaptation practitioners are still thinking overwhelmingly in terms of traditional conservation practices, and have yet to embrace the recommendations from a wide body of conservation literature that climate change necessitates a different emphasis and new approaches.

## Implications for climate funds and project developers

These observations suggest that the vulnerabilities and adaptation needs of other species are not featuring prominently in the adaptation projects developing countries and project developers prioritise for international funding. Further, we notice project proponents significantly overstating the contribution projects are making to ecosystems, suggesting potential benefits for other species that seem disconnected from the project activities and are rarely verified by project monitoring and evaluation (M&E) frameworks. Almost all project documents mention ecosystems or biodiversity, and many claim the project will deliver an indirect benefit for ecosystems – even where there are no substantive activities that seem to address ecosystem health.

To address these gaps in both scale and quality of funding, a number of changes to the way funding is allocated might be considered. First, the gap in funding for ecosystems needs attention at the UNFCCC level, since the Parties to the UNFCCC define the mandate to its climate funds. It is clear that without specific targets or rewards addressing the needs of other species and natural ecosystems, these activities

To turn around these trends, major investments are needed, as is a change in how people prioritise other species in their pursuit of social and economic development

will continue to be marginalised as projects are designed. The UNFCCC could require each of the Funds to set up dedicated funding windows for ecosystem projects, thereby ensuring a greater portion of the available international climate finance goes to projects that specifically improve the resilience of natural ecosystems and the survival prospects of other species. This could also help overcome the fact that most activities to date are small components of projects that target other objectives; hence, the scale of benefits for ecosystems is correspondingly small. There is an urgent need to encourage more large-scale projects, if resilience of other species is to be meaningfully supported.

The results suggest those who are bringing forward funding proposals are under-incentivised to focus on ecosystems in their adaptation agendas; this includes governments and adaptation planners in recipient countries, as well as project developers, including Implementing Entities who are accredited by the Funds to prepare project proposals and help countries implement them. Effort is therefore needed to raise the profile of adaptation for other species, including by helping governments and civil society organisations understand the socio-economic benefits of ensuring the long-term prosperity of ecosystems, as well as the actions needed to ensure this at scale.

While the project documents are prepared by adaptation planners and project developers, the Fund still has an important role to play in dialogue with project proponents and countries as proposals are brought forward. In reviewing funding applications, the Fund could offer guidance on maximising the co-benefits of projects for other species. Their scope for doing this would be enhanced if, again, the UNFCCC provided them with a stronger mandate and/or resources to work with developing countries in the design of projects. Appointing dedicated ecosystems specialists to their technical committees and boards would enable the Fund to more effectively review all proposals to provide expert advice to project developers on the likely impacts and potential benefits of project activities, and would contribute to further elaboration of how ecosystem resilience could be improved. At a minimum, the climate funds should require project developers to include, in their M&E frameworks, indicators relating to changes in ecosystem *quality* when there are ecosystem activities.

Enhanced reporting on these issues would also be welcomed. To this end, it would be helpful if the climate funds – and bilateral donors – made deliberate efforts to report on these questions. That would include not only tagging expenditures against biodiversity objectives (as they do when reporting financial support annually to the OECD's Development Assistance Committee) but also conducting robust, qualitative assessments on whether and how projects are specifically targeting improvements in ecosystem resilience and supporting the adaptation of animals and plants to climate change. The Adaptation Fund, for example, tends to publicly highlight its achievements in terms of the number of beneficiaries but rarely, if ever, mentions what outcomes it has achieved for other species. Though not required under the Fund's current operating guidelines, an evaluation of its accomplishments in light of a wider set of international needs could help further debate about how limited adaptation funding might be prioritised in future.

Finally, these findings might suggest that the conservation community has been slow in interacting with climate finance opportunities, or that there are insufficient opportunities for meaningful involvement. If so, effort is needed to bring in this expertise sooner rather than later.

At the country level, limited funding opportunities may have understandably led to the prioritisation of adaptation projects with a strong socio-economic development angle. However, as argued in Section 1, socio-economic outcomes around the world depend heavily on ecosystem health in the long-term. The way ecosystems are considered in sustainable development agendas might need to be discussed with this longer-term view in mind. Otherwise, we may continue to see a major gap in funding the climate resiliency of other species, and the ecosystems on which they and we depend.

## Concluding remarks

This paper's exploratory analysis of the Adaptation Fund offers only a limited view of what is happening across the landscape of climate funds and bilateral development cooperation on climate adaptation. Further analysis of other funding flows – in other climate funds such as the Green Climate Fund, Global Environment Facility and World Bank's Climate Investment Funds, as well as other biodiversity funding channels – would therefore help to improve our understanding of whether the needs and vulnerabilities of other species are being given priority as the world prepares for the impacts of climate change.

As climate change puts both environments and financial resources under greater stress, we might expect there will be even greater competition and conflict between the needs of humans and of other species. In the contest for limited financial resources, our findings highlight the risk that people may find it easy to neglect the needs and vulnerabilities of other species, and of ecosystems broadly, and instead prioritise funding for interventions that address the physical assets and economic vulnerabilities of people. Our findings seem to confirm a trend, observed in the conservation literature, of a shift away from "naturalness" in conservation and towards the protection or prioritization of ecosystems that are deemed of ecological service on the basis of their contribution to human welfare and enjoyment (Starzomski 2013). If conservation interventions are designed to maximize the benefits nature provides people – meaning species and ecosystems are prioritized for protection on the basis of their assessable utility for humans – this will make life for other species even more precarious.

Addressing the needs and vulnerabilities of other species is urgent. Hannah et al (2011) argue that there is a narrow time window in some places in which populations can persist as the climate changes, and that after a certain time it will become impossible to salvage natural systems or save certain species. In the meantime, continued patterns of ecosystem degradation and land use change will have further limited the options for establishing protected areas or connections between discrete habitat patches.

To turn around these trends, major investments are needed, as is a change in how people prioritise other species in their pursuit of social and economic development. In light of the broad, overwhelming trend of ecosystem devastation globally, small adaptation projects may seem of miniscule importance. Yet the persistent degradation and fragmentation of natural ecosystems makes effective use of adaptation funding even more critical, to ensure the natural world can cope with and adapt to climate change.

## References

- Ackerly, D. D., Loarie, S. R., Cornwell, W. K., Weiss, S. B., Hamilton, H., Branciforte, R. and Kraft, N. J. B. (2010). The geography of climate change: implications for conservation biogeography. *Diversity and Distributions*, 16(3). 476–87. DOI:10.1111/j.1472-4642.2010.00654.x.
- Adaptation Fund (2011). *Results Framework and Baseline Guidance: Project-Level*. Washington, DC. <https://www.adaptation-fund.org/document/results-framework-and-baseline-guidance-project-level>.
- Alagador, D., Cerdeira, J. O. and Araújo, M. B. (2016). Climate change, species range shifts and dispersal corridors: an evaluation of spatial conservation models. *Methods in Ecology and Evolution*, 7(7). 853–66. DOI:10.1111/2041-210X.12524.
- Araujo, M. B., Cabeza, M., Thuiller, W., Hannah, L. and Williams, P. H. (2004). Would climate change drive species out of reserves? An assessment of existing reserve-selection methods. *Global Change Biology*, 10(9). 1618–26. DOI:10.1111/j.1365-2486.2004.00828.x.
- Ashcroft, M. B., Chisholm, L. A. and French, K. O. (2009). Climate change at the landscape scale: predicting fine-grained spatial heterogeneity in warming and potential refugia for vegetation. *Global Change Biology*, 15(3). 656–67. DOI:10.1111/j.1365-2486.2008.01762.x.
- Balmford, A. (2002). Economic reasons for conserving wild nature. *Science*, 297(5583). 950–53. DOI:10.1126/science.1073947.
- Beier, P. and Brost, B. (2010). Use of Land Facets to Plan for Climate Change: Conserving the Arenas, Not the Actors: Land Facets for Climate-Change Planning. *Conservation Biology*, 24(3). 701–10. DOI:10.1111/j.1523-1739.2009.01422.x.
- Berry, P. (2007). *Adaptation Options on Natural Ecosystems: A Report to the UNFCCC Financial and Technical Support Division*. Environmental Change Institute, University of Oxford. [http://unfccc.int/files/cooperation\\_and\\_support/financial\\_mechanism/application/pdf/berry.pdf](http://unfccc.int/files/cooperation_and_support/financial_mechanism/application/pdf/berry.pdf).
- Bonn, A., Macgregor, N., Stadler, J., Korn, H., Stiffel, S., Wolf, K. and van Dijk, N. (2014). *Helping Ecosystems in Europe to Adapt to Climate Change*. BfN-Skripten 375. Bundesamt für Naturschutz (Federal Agency for Nature Conservation), Bonn. [http://www.bfn.de/fileadmin/MDB/documents/service/Skript\\_375.pdf](http://www.bfn.de/fileadmin/MDB/documents/service/Skript_375.pdf).
- Brooke, C. (2008). Conservation and Adaptation to Climate Change. *Conservation Biology*, 22(6). 1471–76. DOI:10.1111/j.1523-1739.2008.01031.x.
- Burns, C. E., Johnston, K. M. and Schmitz, O. J. (2003). Global climate change and mammalian species diversity in U.S. national parks. *Proceedings of the National Academy of Sciences*, 100(20). 11474–77. DOI:10.1073/pnas.1635115100.
- Burrows, M. T., Schoeman, D. S., Richardson, A. J., Molinos, J. G., Hoffmann, A., et al. (2014). Geographical limits to species-range shifts are suggested by climate velocity. *Nature*, 507(7493). 492–95. DOI:10.1038/nature12976.
- Chornesky, E. A., Ackerly, D. D., Beier, P., Davis, F. W., Flint, L. E., et al. (2015). Adapting California's Ecosystems to a Changing Climate. *BioScience*, 65(3). 247–62. DOI:10.1093/biosci/biu233.
- Clews, L. (2012). How Australian legislation can consider climate change in ecological impact assessment. In *Wildlife and Climate Change: towards robust conservation strategies for Australian fauna*. D. Lunney and P. Hutchings (eds.). Royal Zoological Society of New South Wales, New South Wales, Australia. 144–49. <http://publications.rzsnsw.org.au/doi/pdf/10.7882/FS.2012.021?code=rzsw-site>.
- Convention on Biological Diversity (2016). *Synthesis Report on the Experience with Ecosystem-Based Approaches to Climate Change Adaptation and Disaster Risk Reduction*. Convention on Biological Diversity, UNEP, Washington DC. <https://www.cbd.int/doc/meetings/sbstta/sbstta-20/information/sbstta-20-inf-02-en.pdf>.
- Cotton, P. A. (2003). Avian migration phenology and global climate change. *Proceedings of the National Academy of Sciences*, 100(21). 12219–22. DOI:10.1073/pnas.1930548100.
- Cross, M. S., Zavaleta, E. S., Bachelet, D., Brooks, M. L., Enquist, C. A. F., et al. (2012). The Adaptation for Conservation Targets (ACT) Framework: A Tool for Incorporating Climate Change into Natural Resource Management. *Environmental Management*, 50(3). 341–51. DOI:10.1007/s00267-012-9893-7.
- DEFRA (2008). *England Biodiversity Strategy Climate Change Adaptation Principles. Conserving Biodiversity in a Changing Climate*. DEFRA, London. [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/69270/pb13168-ebc-ccap-081203.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/69270/pb13168-ebc-ccap-081203.pdf).
- Department of Environment, Climate Change and Water (2010). *Priorities for Biodiversity Adaptation to Climate Change*. Department of Environment, Climate Change and Water (NSW), Sydney, Australia. <http://www.environment.nsw.gov.au/resources/biodiversity/10771prioritiesbioadaptcc.pdf>.
- Donaldson, L., Wilson, R. J. and Maclean, I. M. D. (2017). Old concepts, new challenges: adapting landscape-scale conservation to the twenty-first century. *Biodiversity and Conservation*, 26(3). 527–52. DOI:10.1007/s10531-016-1257-9.
- Environmental Defenders Office (2009). *Climate Change and the Legal Framework for Biodiversity Protection in Australia: A Legal and Scientific Analysis*. Environmental Defenders Office, Sydney. [http://d3n8a8pro7vnm.cloudfront.net/edonsw/pages/282/attachments/original/1380667846/090724cth\\_discussion\\_paper.pdf?1380667846](http://d3n8a8pro7vnm.cloudfront.net/edonsw/pages/282/attachments/original/1380667846/090724cth_discussion_paper.pdf?1380667846).

- European Commission (2013). *Guidelines on Climate Change and Natura 2000*. 2013–068. European Commission, Brussels. <http://ec.europa.eu/environment/nature/climatechange/pdf/Guidance%20document.pdf>.
- Feger, C. and Pirard, R. (2011). *Assessing Funding Needs for Biodiversity: Critical Issues*. 06/11. Institut du développement durable et des relations internationales (IDDRI), Paris, France. <http://www.iddri.org/Publications/Assessing-funding-needs-for-biodiversity-Critical-issues>.
- Gallagher, R. V., Makinson, R. O., Hogbin, P. M. and Hancock, N. (2015). Assisted colonization as a climate change adaptation tool. *Austral Ecology*, 40(1). 12–20. DOI:10.1111/aec.12163.
- Game, E. T., Lipsett-Moore, G., Saxon, E., Peterson, N. and Sheppard, S. (2011). Incorporating climate change adaptation into national conservation assessments. *Global Change Biology*, 17(10). 3150–60. DOI:10.1111/j.1365-2486.2011.02457.x.
- Gaston, K. J. (2009). Geographic range limits of species. *Proceedings of the Royal Society B: Biological Sciences*, 276(1661). 1391–93. DOI:10.1098/rspb.2009.0100.
- Gimona, A., Poggio, L., Polhill, J. G. and Castellazzi, M. (2015). Habitat networks and food security: promoting species range shift under climate change depends on life history and the dynamics of land use choices. *Landscape Ecology*, 30(5). 771–89. DOI:10.1007/s10980-015-0158-8.
- Glick, P., Chmura, H. and Stein, B. A. (2011). *Moving the Conservation Goalposts: A Review of Climate Change Adaptation Literature*. National Wildlife Federation, Washington D.C.
- Glick, P., Stein, B. A. and Edelson, N. A. (2011). *Scanning the Conservation Horizon: A Guide to Climate Change Vulnerability Assessment*. National Wildlife Federation, Washington, DC. <http://www.nwf.org/~media/pdfs/global-warming/climate-smart-conservation/nwfs scanning the conservation horizon final 92311.ashx>.
- Groves, C. R., Game, E. T., Anderson, M. G., Cross, M., Enquist, C., et al. (2012). Incorporating climate change into systematic conservation planning. *Biodiversity and Conservation*, 21(7). 1651–71. DOI:10.1007/s10531-012-0269-3.
- Gunderson, L. and Holling, C. S. (2002). *Panarchy: Understanding Transformations in Human and Natural Systems*. Island Press, New York.
- Halpin, P. N. (1997). Global Climate Change and Natural-Area Protection: Management Responses and Research Directions. *Ecological Applications*, 7(3). 828. DOI:10.2307/2269436.
- Hannah, L. (2011). Climate Change, Connectivity, and Conservation Success. *Conservation Biology*, 25(6). 1139–42. DOI:10.1111/j.1523-1739.2011.01788.x.
- Hannah, L., Dave, R., Lowry, P. P., Andelman, S., Andrianarisata, M., et al. (2008). Climate change adaptation for conservation in Madagascar. *Biology Letters*, 4(5). 590–94. DOI:10.1098/rsbl.2008.0270.
- Hannah, L. J. (2015). *Climate Change Biology*. Second Edition. Academic Press, Elsevier, Amsterdam.
- Hannah, L., Midgley, G., Andelman, S., Araujo, M., Hughes, G., Martinez-Meyer, E., Pearson, R. and Williams, P. (2007). Protected area needs in a changing climate. *Frontiers in Ecology and the Environment*, 2007(5). 131–38.
- Hannah, L., Midgley, G. F., Lovejoy, T., Bond, W. J., Bush, M., Lovett, J. C., Scott, D. and Woodward, F. I. (2002). Conservation of Biodiversity in a Changing Climate. *Conservation Biology*, 16(1). 264–68. DOI:10.1046/j.1523-1739.2002.00465.x.
- Harris, J. A., Hobbs, R. J., Higgs, E. and Aronson, J. (2006). Ecological Restoration and Global Climate Change. *Restoration Ecology*, 14(2). 170–76. DOI:10.1111/j.1526-100X.2006.00136.x.
- Harris, R. M. B., Carter, O., Gilfedder, L., Porfirio, L. L., Lee, G. and Bindoff, N. L. (2015). Noah's Ark Conservation Will Not Preserve Threatened Ecological Communities under Climate Change. *PLOS ONE*, 10(4). e0124014. DOI:10.1371/journal.pone.0124014.
- Heller, N. E., Kreidler, J., Ackerly, D. D., Weiss, S. B., Recinos, A., Branciforte, R., Flint, L. E., Flint, A. L. and Micheli, E. (2015). Targeting climate diversity in conservation planning to build resilience to climate change. *Ecosphere*, 6(4). art65. DOI:10.1890/ES14-00313.1.
- Heller, N. E. and Zavaleta, E. S. (2009). Biodiversity management in the face of climate change: A review of 22 years of recommendations. *Biological Conservation*, 142(1). 14–32. DOI:10.1016/j.biocon.2008.10.006.
- Hodgson, J. A., Thomas, C. D., Wintle, B. A. and Moilanen, A. (2009). Climate change, connectivity and conservation decision making: back to basics. *Journal of Applied Ecology*, 46(5). 964–69. DOI:10.1111/j.1365-2664.2009.01695.x.
- Hoegh-Guldberg, O., Hughes, L., McIntyre, S., Lindenmayer, D. B., Parmesan, C., Possingham, H. P. and Thomas, C. D. (2008). Assisted Colonization and Rapid Climate Change. *Science*, 321(5887). 345–46. DOI:10.1126/science.1157897.
- Hole, D. G., Huntley, B., Arinaitwe, J., Butchart, S. H. M., Collingham, Y. C., Fishpool, L. D. C., Pain, D. J. and Willis, S. G. (2011). Toward a Management Framework for Networks of Protected Areas in the Face of Climate Change. *Conservation Biology*, 25(2). 305–15. DOI:10.1111/j.1523-1739.2010.01633.x.
- Holling, C. S. (1973). Resilience and stability of ecological systems. *Annual review of ecology and systematics*, 4. 1–23. <http://www.jstor.org/stable/2096802>.

- Hopper, S. D. (2007). New Life for Systematics. *Science*, 316(5828). 1097–1097. DOI:10.1126/science.1144898.
- Hulme, P. E. (2005). Adapting to climate change: is there scope for ecological management in the face of a global threat? *Journal of Applied Ecology*, 42(5). 784–94. DOI:10.1111/j.1365-2664.2005.01082.x.
- Hunter Jr., M., Dinerstein, E., Hoekstra, J. and Lindenmayer, D. (2010). A Call to Action for Conserving Biological Diversity in the Face of Climate Change: Editorial. *Conservation Biology*, 24(5). 1169–71. DOI:10.1111/j.1523-1739.2010.01569.x.
- Johnson, J. E. and Welch, D. J. (2009). Marine Fisheries Management in a Changing Climate: A Review of Vulnerability and Future Options. *Reviews in Fisheries Science*, 18(1). 106–24. DOI:10.1080/10641260903434557.
- Johnson, K. H., Vogt, K. A., Clark, H. J., Schmitz, O. J. and Vogt, D. J. (1996). Biodiversity and the productivity and stability of ecosystems. *Trends in Ecology & Evolution*, 11(9). 372–77. DOI:10.1016/0169-5347(96)10040-9.
- Jones, K. R., Watson, J. E. M., Possingham, H. P. and Klein, C. J. (2016). Incorporating climate change into spatial conservation prioritisation: A review. *Biological Conservation*, 194. 121–30. DOI:10.1016/j.biocon.2015.12.008.
- Jump, A. S. and Penuelas, J. (2005). Running to stand still: adaptation and the response of plants to rapid climate change. *Ecology Letters*, 8(9). 1010–20. DOI:10.1111/j.1461-0248.2005.00796.x.
- Kareiva, P., Enquist, C., Johnson, A., Julius, S., Lawler, J., Petersen, B., Pitelka, L., Shaw, R. and West, J. (2008). Synthesis and conclusions. In *Preliminary review of adaptation options for climate-sensitive ecosystems and resources. A report by the U.S Climate Change Science Program and the Subcommittee on Global Change Research*. S. Julius and J. West (eds.). U.S EPA, Washington, DC. <https://pubs.er.usgs.gov/publication/70182094>.
- Keppel, G., Van Niel, K. P., Wardell-Johnson, G. W., Yates, C. J., Byrne, M., Mucina, L., Schut, A. G. T., Hopper, S. D. and Franklin, S. E. (2012). Refugia: identifying and understanding safe havens for biodiversity under climate change. *Global Ecology and Biogeography*, 21(4). 393–404. DOI:10.1111/j.1466-8238.2011.00686.x.
- Krosby, M., Tewksbury, J., Haddad, N. M. and Hoekstra, J. (2010). Ecological Connectivity for a Changing Climate: Ecological Connectivity. *Conservation Biology*, 24(6). 1686–89. DOI:10.1111/j.1523-1739.2010.01585.x.
- Lawler, J., Watson, J. and Game, E. (2015). Conservation in the face of climate change: recent developments. *F1000Research*, 4(F1000 Faculty Rev). 1158. DOI:10.12688/f1000research.6490.1.
- Lawton, J. H., Brotherton, P. N. M., Brown, V. K., Elphick, C., Fitter, A. H., et al. (2010). *Making Space for Nature: A Review of England's Wildlife Sites and Ecological Network*. DEFRA, London. <http://webarchive.nationalarchives.gov.uk/20101109170125/http://www.defra.gov.uk/environment/biodiversity/documents/201009space-for-nature.pdf>.
- Lindenmayer, D., Hobbs, R. J., Montague-Drake, R., Alexandra, J., Bennett, A., et al. (2007). A checklist for ecological management of landscapes for conservation. *Ecology Letters*, 11(1). 78–91. DOI:10.1111/j.1461-0248.2007.01114.x.
- Lunt, I. D., Byrne, M., Hellmann, J. J., Mitchell, N. J., Garnett, S. T., et al. (2013). Using assisted colonisation to conserve biodiversity and restore ecosystem function under climate change. *Biological Conservation*, 157. 172–77. DOI:10.1016/j.biocon.2012.08.034.
- Maciver, D. C. and Wheaton, E. (2005). Tomorrow's Forests: Adapting to A Changing Climate. *Climatic Change*, 70(1–2). 273–82. DOI:10.1007/s10584-005-5950-z.
- Martin, S. (2011). Ecosystem-Based Adaptation: What Does It Really Mean? *WWF Climate Prep: Adaptation Stories*. <http://www.climateprep.org/stories/2016/3/1/ecosystem-based-adaptation-what-does-it-really-mean>.
- Mawdsley, J. R., O'Malley, R. and Ojima, D. S. (2009). A Review of Climate-Change Adaptation Strategies for Wildlife Management and Biodiversity Conservation. *Conservation Biology*, 23(5). 1080–89. DOI:10.1111/j.1523-1739.2009.01264.x.
- McCarthy, D. P., Donald, P. F., Scharlemann, J. P. W., Buchanan, G. M., Balmford, A., et al. (2012). Financial Costs of Meeting Global Biodiversity Conservation Targets: Current Spending and Unmet Needs. *Science*, 338(6109). 946–49. DOI:10.1126/science.1229803.
- McClanahan, T. R., Cinner, J. E., Maina, J., Graham, N. A. J., Daw, T. M., et al. (2008). Conservation action in a changing climate. *Conservation Letters*, 1(2). 53–59. DOI:10.1111/j.1755-263X.2008.00008\_1.x.
- Millar, C. I., Stephenson, N. L. and Stephens, S. L. (2007). Climate Change and Forests of the Future: Managing in the Face of Uncertainty. *Ecological Applications*, 17(8). 2145–51. DOI:10.1890/06-1715.1.
- Millennium Ecosystem Assessment (Program), ed. (2005). *Ecosystems and Human Well-Being: Synthesis*. Island Press, Washington, DC.
- Morecroft, M. D., Crick, H. Q. P., Duffield, S. J. and Macgregor, N. A. (2012). Resilience to climate change: translating principles into practice. *Journal of Applied Ecology*, 49(3). 547–51. DOI:10.1111/j.1365-2664.2012.02136.x.
- Noss, R. (2001). Beyond Kyoto: Forest Management in a Time of Rapid Climate Change. *Conservation Biology*, 15(3). 578–90.



- Núñez, T. A., Lawler, J. J., Mcrae, B. H., Pierce, D. J., Krosby, M. B., Kavanagh, D. M., Singleton, P. H. and Tewksbury, J. J. (2013). Connectivity Planning to Address Climate Change. *Conservation Biology*, 27(2). 407–16. DOI:10.1111/cobi.12014.
- OECD (2003). *Emerging Systemic Risks in the 21st Century: An Agenda for Action*. Organisation for Economic Co-operation and Development, Paris. <http://www.oecd.org/futures/globalprospects/37944611.pdf>.
- Oliver, T. H., Smithers, R. J., Bailey, S., Walmsley, C. A. and Watts, K. (2012). A decision framework for considering climate change adaptation in biodiversity conservation planning. *Journal of Applied Ecology*, 49(6). 1247–55. DOI:10.1111/1365-2664.12003.
- Otero, J., L'Abée-Lund, J. H., Castro-Santos, T., Leonardsson, K., Storch, G. O., et al. (2014). Basin-scale phenology and effects of climate variability on global timing of initial seaward migration of Atlantic salmon (*Salmo salar*). *Global Change Biology*, 20(1). 61–75. DOI:10.1111/gcb.12363.
- Pacifici, M., Foden, W. B., Visconti, P., Watson, J. E. M., Butchart, S. H. M., et al. (2015). Assessing species vulnerability to climate change. *Nature Climate Change*, 5(3). 215–24. DOI:10.1038/nclimate2448.
- Parmesan, C. and Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421. 37–42.
- Pearson, R. G. and Dawson, T. P. (2005). Long-distance plant dispersal and habitat fragmentation: identifying conservation targets for spatial landscape planning under climate change. *Biological Conservation*, 123(3). 389–401. DOI:10.1016/j.biocon.2004.12.006.
- Poiani, K. A., Goldman, R. L., Hobson, J., Hoekstra, J. M. and Nelson, K. S. (2011). Redesigning biodiversity conservation projects for climate change: examples from the field. *Biodiversity and Conservation*, 20(1). 185–201. DOI:10.1007/s10531-010-9954-2.
- Richerzhagen, C., Rodriguez, J. C. and Stepping, K. (2016). *Why We Need More and Better Biodiversity Aid*. 13/2016. German Development Institute (DIE). <https://www.die-gdi.de/en/briefing-paper/article/why-we-need-more-and-better-biodiversity-aid/>.
- Root, T. L., Price, J. T., Hall, K. R., Schneider, S. H., Rosenzweig, C. and Pounds, J. A. (2003). Fingerprints of global warming on wild animals and plants. *Nature*, 421(6918). 57–60. DOI:10.1038/nature01333.
- Rüter, S., Vos, C. C., van Eupen, M. and Rühmkorf, H. (2014). Transboundary ecological networks as an adaptation strategy to climate change: The example of the Dutch – German border. *Basic and Applied Ecology*, 15(8). 639–50. DOI:10.1016/j.baae.2014.09.007.
- Sala, O. E., Chapin, F. S., Armesto, J. J., Berlow, E., Bloomfield, J., et al. (2000). Global Biodiversity Scenarios for the Year 2100. *Science*, 287(5459). 1770–74. DOI:10.1126/science.287.5459.1770.
- Schmitz, O. J., Lawler, J. J., Beier, P., Groves, C., Knight, G., et al. (2015). Conserving Biodiversity: Practical Guidance about Climate Change Adaptation Approaches in Support of Land-use Planning. *Natural Areas Journal*, 35(1). 190–203. DOI:10.3375/043.035.0120.
- Scott, D., Malcolm, J. and Lemieux, C. (2002). Climate Change and Modelled Biome Representation in Canada's National Park System: Implications for System Planning and Park Mandates. *Global Ecology and Biogeography*, 11(6). 475–84.
- Secretariat of the Convention on Biological Diversity (2012). Aichi Biodiversity Targets. <https://www.cbd.int/sp/targets/default.shtml>.
- Settele, J., Scholes, R., Betts, R., Bunn, S. E., Leadley, P., Nepstad, D., Overpeck, J. T. and Taboada, M. A. (2014). Terrestrial and inland water systems. In *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change*. C. B. Field, V. R. Barros, D. J. Dokken, K. J. Mach, M. D. Mastrandrea, et al. (eds.). Cambridge University Press, Cambridge, UK, and New York. 271–359.
- Shafer, C. L. (1999). National park and reserve planning to protect biological diversity: some basic elements. *Landscape and Urban Planning*, 44(2–3). 123–53. DOI:10.1016/S0169-2046(98)00115-7.
- Shoo, L. P., Hoffmann, A. A., Garnett, S., Pressey, R. L., Williams, Y. M., et al. (2013). Making decisions to conserve species under climate change. *Climatic Change*, 119(2). 239–46. DOI:10.1007/s10584-013-0699-2.
- Singh, N. J., Börger, L., Dettki, H., Bunnefeld, N. and Ericsson, G. (2012). From migration to nomadism: movement variability in a northern ungulate across its latitudinal range. *Ecological Applications*, 22(7). 2007–20. DOI:10.1890/12-0245.1.
- Starzomski, B. M. (2013). Novel Ecosystems and Climate Change. In *Novel Ecosystems*. R. J. Hobbs, E. S. Higgs, and C. M. Hall (eds.). John Wiley & Sons, Ltd, Chichester, UK. 88–101. DOI:10.1002/9781118354186.ch10.
- Stein, B. A., Staudt, A., Cross, M. S., Dubois, N. S., Enquist, C., et al. (2013). Preparing for and managing change: climate adaptation for biodiversity and ecosystems. *Frontiers in Ecology and the Environment*, 11(9). 502–10. DOI:10.1890/120277.
- Theobald, D. M., Harrison-Atlas, D., Monahan, W. B. and Albano, C. M. (2015). Ecologically-Relevant Maps of Landforms and Physiographic Diversity for Climate Adaptation Planning. *PLOS ONE*, 10(12). e0143619. DOI:10.1371/journal.pone.0143619.
- Thomas, C. D. (2011). Translocation of species, climate change, and the end of trying to recreate past ecological communities. *Trends in Ecology & Evolution*, 26(5). 216–21. DOI:10.1016/j.tree.2011.02.006.
- Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., et al. (2004). Extinction risk from climate change. *Nature*, 427(6970). 145–48. DOI:10.1038/nature02121.

- Tingley, M. W., Darling, E. S. and Wilcove, D. S. (2014). Fine- and coarse-filter conservation strategies in a time of climate change. *Annals of the New York Academy of Sciences*, 1322(1). 92–109. DOI:10.1111/nyas.12484.
- Tischendorf, L. and Fahrig, L. (2000). How should we measure landscape connectivity? *Landscape Ecology*, 2000(15). 633.
- Townsend, P. and Masters, K. (2015). Lattice-work corridors for climate change: a conceptual framework for biodiversity conservation and social-ecological resilience in a tropical elevational gradient. *Ecology and Society*, 20(2). DOI: 10.5751/ES-07324-200201.
- UN General Assembly (2015). *Transforming Our World: The 2030 Agenda for Sustainable Development*. United Nations, New York.
- UNCBD (2014). *Decision Adopted by the Conference of the Parties for the Convention on Biological Diversity XII/3 Resource Mobilization*. UNEP/CBD/COP/DEC/XII/3. Korea. <https://www.cbd.int/decision/cop/default.shtml?id=13366>.
- UNCBD (2016). *Decision Adopted by the Conference of the Parties to the Convention on Biological Diversity XIII/4. Biodiversity and Climate Change*. CBD/COP/ DEC/XIII/4. Mexico. <https://www.cbd.int/doc/decisions/cop-13/cop-13-dec-04-en.pdf>.
- United Nations (1992a). *Convention on Biological Diversity*. United Nations, Rio de Janeiro. <https://www.cbd.int/doc/legal/cbd-en.pdf>.
- United Nations (1992b). *United Nations Framework Convention on Climate Change*. FCCC/INFORMAL/84. United Nations, New York. <https://unfccc.int/resource/docs/convkp/conveng.pdf>.
- Urban, M. C. (2015). Accelerating extinction risk from climate change. *Science*, 348(6234). 571–73. DOI:10.1126/science.aaa4984.
- van Bodegom, P. M., Verboom, J., Witte, J. P. M., Vos, C. C., Bartholomeus, R. P., Geertsema, W., Cormont, A., van der Veen, M. and Aerts, R. (2013). Synthesis of ecosystem vulnerability to climate change in the Netherlands shows the need to consider environmental fluctuations in adaptation measures. *Regional Environmental Change*, 14(3). 933–42. DOI:10.1007/s10113-013-0511-x.
- van Oppen, M. J. H., Oliver, J. K., Putnam, H. M. and Gates, R. D. (2015). Building coral reef resilience through assisted evolution. *Proceedings of the National Academy of Sciences*, 112(8). 2307–13. DOI:10.1073/pnas.1422301112.
- Vos, C. C., Berry, P., Opdam, P., Baveco, H., Nijhof, B., O'Hanley, J., Bell, C. and Kuipers, H. (2008). Adapting landscapes to climate change: examples of climate-proof ecosystem networks and priority adaptation zones. *Journal of Applied Ecology*, 45(6). 1722–31. DOI:10.1111/j.1365-2664.2008.01569.x.
- Vos, C. C., van der Hoek, D. C. J. and Vonk, M. (2010). Spatial planning of a climate adaptation zone for wetland ecosystems. *Landscape Ecology*, 25(10). 1465–77. DOI:10.1007/s10980-010-9535-5.
- West, J. M., Julius, S. H., Kareiva, P., Enquist, C., Lawler, J. J., Petersen, B., Johnson, A. E. and Shaw, M. R. (2009). U.S. Natural Resources and Climate Change: Concepts and Approaches for Management Adaptation. *Environmental Management*, 44(6). 1001–21. DOI:10.1007/s00267-009-9345-1.
- Wilson, E. and Piper, J. (2008). Spatial planning for biodiversity in Europe's changing climate. *European Environment*, 18(3). 135–51. DOI:10.1002/eet.476.



---

## SEI Headquarters

Linnégatan 87D Box 24218  
104 51 Stockholm Sweden  
Tel: +46 8 30 80 44  
info@sei.org

---

### Måns Nilsson

Executive Director

---

## SEI Africa

World Agroforestry Centre  
United Nations Avenue  
Gigiri P.O. Box 30677  
Nairobi 00100 Kenya  
Tel: +254 20 722 4886  
info-Africa@sei.org

---

### Philip Osano

Centre Director

---

## SEI Asia

15th Floor Witthayakit Building  
254 Chulalongkorn University  
Chulalongkorn Soi 64 Phayathai Road  
Pathumwan Bangkok 10330 Thailand  
Tel: +66 2 251 4415  
info-Asia@sei.org

---

### Niall O'Connor

Centre Director

---

## SEI Tallinn

Arsenal Centre  
Erika 14, 10416  
Tallinn, Estonia  
info-Tallinn@sei.org

---

### Lauri Tammiste

Centre Director

---

## SEI Oxford

Florence House 29 Grove Street  
Summertown Oxford  
OX2 7JT UK  
Tel: +44 1865 42 6316  
info-Oxford@sei.org

---

### Ruth Butterfield

Centre Director

---

---

## SEI US

### Main Office

11 Curtis Avenue  
Somerville MA 02144-1224 USA  
Tel: +1 617 627 3786  
info-US@sei.org

---

### Michael Lazarus

Centre Director

---

## SEI US

### Davis Office

400 F Street  
Davis CA 95616 USA  
Tel: +1 530 753 3035

---

## SEI US

### Seattle Office

1402 Third Avenue Suite 900  
Seattle WA 98101 USA  
Tel: +1 206 547 4000

---

## SEI York

University of York  
Heslington York  
YO10 5DD UK  
Tel: +44 1904 32 2897  
info-York@sei.org

---

### Lisa Emberson

Centre Director

---

## SEI Latin America

Calle 71 # 11-10  
Oficina 801  
Bogota Colombia  
Tel: +57 1 6355319  
info-LatinAmerica@sei.org

---

### David Purkey

Centre Director